


**RHODES UNIVERSITY**  
*Where leaders learn*

# RU PICTOGRAMS

Dosage and frequency




Take once a day




Take in the morning




Take at night



Take morning and night



Take 3 times a day



Take 4 times a day


## Administering and using medicines




In the ear




In the eye




In the nose



In the vagina





In the rectum



Mouthwash – swill and spit




Rub lotion on skin –  
do not swallow




Oral drops –  
swirl in mouth and swallow

## Additional instructions for taking medicines



Take medicine with food



Take medicine on an empty stomach




Do not drink alcohol while taking this medicine



Do not have dairy products with this medicine



Do not take tablets if breastfeeding



Do not take tablets if pregnant




Do not share tablets




Do not share capsules




Take medicine at the correct time




Do not break or crush tablets or capsules




Shake bottle before use




Take tablets with a glass of water



Complete the course - tablets



Complete the course - liquids



Crush tablet and mix with juice, yoghurt, jam, water


## Side effects or indications: pg 1



Constipation



Diarrhoea



Cough



Headache



Dizziness



Fever



Vomiting




Drowsiness



Heartburn



Rash



Full body  
rash



General body pain



Feeling weak and  
tired



Back pain




Stomach ache

## Side effects or indications: pg 2



Loss of appetite



Weight loss




Rash and fever



Hearing problems




Confusion




Peripheral neuropathy



Night sweats



Emaciated and ill



Blurred vision



Nightmares




Yellowing of eyes

## Storage of medicines



Out of reach of children



In a cool dry place



In the fridge




Not in the bathroom



Not near the fire




Not in a hot car



Not in the sun



Not near the stove  
(paraffin)



Not on windowsill  
in the sun




In a packet hung high  
on the wall – if no  
other safe place  
available




## Tablets, capsules, bottles, droppers




Bottle



Bottle and dropper




Tablets and capsules




Tablets-shape 1



Tablets-shape 2



Tablets-shape 3




Capsules




## Miscellaneous



Friends hugging



Shaking hands



Open window for ventilation




Queue for toilet



Wear a mask



Thumbs up – feeling good



Sharing food



Sangoma-traditional healer

**Checkers  
Shoprite  
Pick 'n Pay**

Supermarkets

**⊕  
PHARMACY**


Pharmacy

**⊕  
CLINIC**

Clinic



Informal shop




No smoking


# TB Pictograms




Cough-good safe practice



Cough+droplet spread – bad practice




Sneeze+droplet spread – bad practice




Spitting near others- bad practice




Coughing up blood



Open window for ventilation




2 weeks of coughing - calendar



Sputum bottle



Sputum collection sequence



Clearing of lungs with treatment

## TB Pictograms

These depictions of the TB bacterium were used in creating a 'story' about sensitive and drug-resistant TB

Sensitive TB bacterium -  
alive




Sensitive TB bacterium -  
dead



Drug-resistant TB  
bacterium – strong +  
dangerous



Drug-resistant TB  
bacterium - dead



TB bacteria-group

